

# Management Of Rotator Cuff Tears In Overhead Athletes

**Brian Serrano<sup>1,2</sup>, D.C., Ph.D., ATC, CSCS, TSAC-F, OPE-SC, CES**

<sup>1</sup>Director of Rehabilitation and Performance, HPI Sports Medicine and Performance

<sup>2</sup>Assistant Professor, University of Medical Science in Arizona

Corresponding Author: Dr. Brian Serrano

28062 Forbes Road, Unit D

Laguna Niguel, CA 92677

[Brianserrano171@gmail.com](mailto:Brianserrano171@gmail.com)

818-926-7269

## Abstract

**Introduction:** The rotator cuff is made of 4 muscles that may be commonly be injured in overhead sports. Tears to the rotator cuff are common pathologies that are difficult to treat and to date there are no clinical guidelines or best practices for treating tears in the athletic population. The purpose of this mini review article is to synthesize the best evidence within the last 5 years in the management of rotator cuff tears in the overhead athlete.

**Materials and Methods:** A literature review was performed using terms to include shoulder injuries, rotator cuff injuries, and overhead athletes in multiple validated databases from the years 2015-2020.

**Results:** The initial query resulted in 8,852 articles available for analysis. After the initial analysis for duplicates and full text 6,000 articles were left. The search for relevant article that met the inclusion yielded 100 articles to be included into the final review. These articles were instrumental in synthesizing the various treatment and management of rotator cuff tears.

**Discussion:** In the overhead athlete, the first line of defense should be conservative care for 6-8 weeks to see if patient reported outcomes can be improved. Advanced imaging in the form of MRI should be obtained to assess the degree of damage to the rotator cuff. If the athlete is in season, then orthobiologics may be explored as an option. However, if the athlete does not achieve pre-injury levels of function then arthroscopic surgery may be indicated for optimal performance.

**Conclusion:** Rotator cuff tears are common in the overhead athletes, and currently there are no guidelines in their treatment and management. This mini review found evidence to support a conservative approach, followed by a possible orthobiologic intervention, and finally arthroscopic surgery if the athlete wishes to return to performance.

## Introduction

Rotator cuff injury is common in overhead athletes in both throwing and non-throwing sports<sup>1-3</sup>. In throwing sports, the throwing motion imparts a large amount of stress in the shoulder joint<sup>4,5</sup>. The humeral

angular velocities can reach from 7000-8000 degrees/second<sup>6,7</sup>. The shoulder joint (Glenohumeral joint) is a ball and socket joint that is inherently mobile. The stability of the shoulder can be categorized as static and dynamic. Static stabilizers include the glenohumeral ligaments, joint capsule, and labrum<sup>8-10</sup>. The dynamic stabilizers are made up of the rotator cuff musculature which serve to center the humeral head within the glenoid cavity<sup>11,12</sup>. The rotator cuff consists of the: supraspinatus, infraspinatus, teres minor, and subscapularis<sup>13,14</sup>. These muscles serve to resist distraction, horizontal adduction, and internal rotation<sup>15-17</sup>. During the deceleration phase of the throwing phase, the posterior rotator cuff must work eccentrically to slow down the distraction forces produced at the shoulder<sup>18</sup>. In non-throwing sports such as Olympic Weightlifting, the turnover phase of the Clean and Snatch produces large amounts of torque at the shoulder<sup>19</sup>. During the Jerk, the body must pre-load the shoulder through a short dip in the lower extremity before the barbell is lifted overhead<sup>20</sup>. If the transition from the lower body to the upper extremity is not efficient this may cause a concentration of load directly onto the shoulder and rotator cuff<sup>20-22</sup>. It is during these motions that injury can occur. More importantly, due to volume and repetition performed by overhead athletes, rotator cuff injury may more often be chronic or degenerative<sup>23,24</sup>. These chronic or degenerative injuries may lead to partial thickness or full thickness tears. In a partial thickness tear only part of tendon has torn off the bone while the entire tendon has separated or torn from the bone in a full thickness tear<sup>25-28</sup>. Tears to the rotator cuff have been investigated thoroughly in the literature, however no treatment and management guidelines seem to exist. The purpose of this mini-review is to synthesize the best evidence published in the last 5 years to come up with guidelines for treatment and management of rotator cuff tears in overhead athletes.

## Methods

A literature search was performed on PubMed with the terms "rotator cuff tear", "rotator cuff injury", "rotator cuff pathology", "overhead athlete", and "shoulder injuries" with a publication date from 2015-2020 to capture the most recent data on this topic.

Upon the initial review was complete, the author (BS) went through an initial screening of the results using the title to ensure articles were not duplicates and date of publication was in the inclusion criteria. After this initial screen, all articles of interest were reviewed using a combination of the abstract and full text article. If the author (BS) had any doubts or questions about including an article, the corresponding author would try to be reached. For final inclusion into the mini review, the article had to meet all criteria set forth above.

## Results

The initial query resulted in 8,852 articles available for analysis. After the initial analysis for duplicates and full text 6,000 articles were left. The search for relevant article that met the inclusion yielded 100 articles to be included into the final review. These articles were instrumental in synthesizing the various treatment and management of rotator cuff tears. Specifically, articles included definitions of partial thickness vs full thickness tears, conservative management, orthobiologics, and arthroscopic surgery.

## Discussion

Rotator cuff tears are common in the overhead athletes, with chronic and degenerative tears having a higher incidence rate<sup>29</sup>. The supraspinatus and infraspinatus are the most commonly affected muscles<sup>30,31</sup>. Rotator cuff tears may include various signs and symptoms which include generalized pain the shoulder, referred pain to the lateral shoulder, pain with sudden movements, loss of range of motion, and loss of strength<sup>32-34</sup>. When performing the physical exam in the overhead athletes it is important rule out the cervical spine as the source of injury. Furthermore, the proximal biceps and labrum are commonly injured within this population. SLAP tears are the most common<sup>35,36</sup>, however because the labrum is a 360 degree structure the clinician must be conscious on injuries to the inferior labrum (Bankart lesion)<sup>37</sup> and posterior labrum (Kim's lesion)<sup>38</sup>. The specific guidelines for treatment will depend on level of play, stage of season, and the amount of limitation the athlete is experiencing. For example, partial thickness tears respond well to conservative treatment in short-term and long-term follow-up. The athlete, clinician, coach, and orthopedic surgeon must all be informed to the progression of the tear into a larger tear in the future.

Diagnosis of a rotator cuff tear can be performed in many ways such as clinically or with imaging. Using diagnostic ultrasound, CT scan, and MRI all provide well validated avenues of advanced imaging. MRI does remain as the gold standard due to its high sensitivity and specificity<sup>39-41</sup>. Diagnostic ultrasound is reliable but is highly dependent on user expertise and familiarity<sup>42</sup>. CT scan presents a quicker way than MRI but the high amounts of radiation and lower sensitivity to soft tissue injuries make it less used<sup>43</sup>.

Conservative treatment is the first line of defense once a rotator cuff tear is suspected or diagnosed. Best guidelines for conservative treatment include the athlete as a holistic body using the concept of kinetic chain and regional interdependence. This means that even though the shoulder is injured, the clinician should assess the entire body with emphasis on scapular neuromuscular control, thoracic spine mobility, the hip (femoroacetabular joint), and ankle joint (talocrural joint)<sup>44,45</sup>. The literature shows that athletes with shoulder injuries have poor neuromuscular control in that the scapula can't achieve retraction, adduction, and posterior tilt<sup>46</sup>. In the thoracic spine, if extension can't be achieved properly, the shoulder must compensate and put more stress on local structures like the rotator cuff and labrum<sup>47</sup>. The hip must have adequate single-leg stability if the athlete is throwing and in general must have adequate internal rotation to achieve motion and avoid stress in the lumbar spine setting of a chain reaction as energy is transferred into the upper extremity<sup>48</sup>. Similarly, the ankle joint must have adequate dorsiflexion for efficient transferring of force into the trunk and upper extremity. Conservative treatment should be attempted consecutively for 6-8 before exploring other options.

Recently, the use of orthobiologics has increased dramatically in sports medicine due to its low morbidity, low levels of invasiveness, and a quicker return to play<sup>49</sup>. Although there are many definitions of what an orthobiologic is, this review will focus on platelet rich plasma (PRP) and Stem Cell Therapy. It is interesting to note that many arthroscopic procedures augment their repair using PRP but this section will only summarize findings using PRP as a standalone intervention. PRP is an autologous platelet concentrate that typically contains more than 1000 x 10<sup>3</sup> platelets/ML, representing a three to five fold increase as compared to whole blood. PRP contains more than 1500 bioactive growth factors that promote tendon healing<sup>50,51</sup>. It is theorized to have a mechanism of action through modulation of the inflammatory pathway which encourages healing of tendon, ligament, muscle, and bone<sup>52-54</sup>. The reader should be aware of the two main variations (leukocyte-rich and leukocyte-poor preparations) of which there is no clear superior option at the time of writing this review. Shams et al. found evidence for PRP injection as a good alternative to corticosteroid injection for patients with partial rotator cuff tears<sup>55</sup>. Conversely, Kesikburns found no difference in pain or functional outcome scores at any time point up to 1 year between PRP and placebo injections<sup>55</sup>.

Stem Cell therapy (SCT) consists of mesenchymal stem cells (MSC) which differentiate into different mesodermal tissues and have strong anti-inflammatory, immunoregulatory, and angiogenic potential<sup>56,57</sup>. Thus, SCT is seen as a potentially effective therapy in the healing of rotator cuff tears. SCT has been investigated in the healing of the tendon-bone interface which is crucial in regaining

biomechanical strength after injury<sup>58-60</sup>. The most common source for SCT is autologous bone marrow while other less commonly used sources include adipose tissue derived MSC's<sup>61</sup>. The biggest obstacle with SCT is the cost which is not covered by insurance and the lack of published studies in humans. To date, there is no consensus statement on the use of SCT, it is an experimental treatment that seems to show promising results.

Surgical management of rotator cuff tears should be viewed as a necessary last resort if the athlete has not achieved their necessary function for optimal performance. Surgically, there are three common procedures performed: open, mini-open, and arthroscopic. Arthroscopic has become the dominant technique in the last 10 years due to its lower level of invasiveness, faster return to play time, and higher patient reported outcomes<sup>62,63</sup>. Needless to say, the rehabilitation period after arthroscopic is longer as compared to conservative treatment, PRP, or SCT and involves sling use for about 4-6 weeks, active range of motion starting at 8 weeks, strengthening exercises at week 10<sup>64-66</sup>. The athlete may begin their return to gradual progression starting at Month 4 (20 weeks) depending on their sport. In the throwing athlete, the throwing progression is usually started at month 6 (week 24)<sup>62,63,66</sup>.

### Conclusion

Rotator cuff tears are common in the overhead athlete population with chronic/degenerative tears being more prevalent compared to acute tears. The general treatment and management will depend on competition level, time of season, and symptom interference in the athlete. The purpose of this mini-review was to synthesize the latest research and synthesize it into recommendations for the clinician working with the overhead population. Upon clinical or imaging diagnosis, conservative treatment should be attempted for 6-8 weeks. If this is not sufficient, a physician should be consulted to explore options of PRP or SCT. If this option is explored or bypassed, the last step would be arthroscopic surgery with an orthopedic surgeon, preferably one who is fellowship trained in sports medicine.

### References

1. Page P. Shoulder muscle imbalance and subacromial impingement syndrome in overhead athletes. *International journal of sports physical therapy*. 2011;6(1):51-58. Accessed August 23, 2020. <http://www.ncbi.nlm.nih.gov/pubmed/21655457>
2. Burkhart SS, Morgan CD, Kibler WB. Shoulder injuries in overhead athletes: The "dead arm" revisited. *Clinics in Sports Medicine*. 2000;19(1):125-158. doi:10.1016/S0278-5919(05)70300-8
3. Brockmeier SF, Dodson CC, Gamradt SC, Coleman SH, Altchek DW. Arthroscopic
4. Gainor BJ, Piotrowski G, Puhl J, Allen WC, Hagen R. The throw: Biomechanics and acute injury. *The American Journal of Sports Medicine*. 1980;8(2):114-118. doi:10.1177/036354658000800210
5. Burkhart SS, Morgan CD, ben Kibler W. The disabled throwing shoulder: Spectrum of pathology Part I: Pathoanatomy and biomechanics. *Arthroscopy - Journal of Arthroscopic and Related Surgery*. 2003;19(4):404-420. doi:10.1053/jars.2003.50128
6. Werner SL, Gill TJ, Murray TA, Cook TD, Hawkins RJ. Relationships between throwing mechanics and shoulder distraction in professional baseball pitchers. *American Journal of Sports Medicine*. 2001;29(3):354-358. doi:10.1177/03635465010290031701
7. Werner SL, Suri M, Guido JA, Meister K, Jones DG. Relationships between ball velocity and throwing mechanics in collegiate baseball pitchers. *Journal of Shoulder and Elbow Surgery*. 2008;17(6):905-908. doi:10.1016/j.jse.2008.04.002
8. Rodosky MW, Harner CD, Fu FH. The Role of the Long Head of the Biceps Muscle and Superior Glenoid Labrum in Anterior Stability of the Shoulder. *The American Journal of Sports Medicine*. 1994;22(1):121-130. doi:10.1177/036354659402200119
9. Ovesen J, Nielsen S. Stability of the shoulder joint: Cadaver study of stabilizing structures. *Acta Orthopaedica*. 1985;56(2):149-151. doi:10.3109/17453678508994342
10. O'Connell PW, Nuber GW, Mileski RA, Lautenschlager E. The contribution of the glenohumeral ligaments to anterior stability of the shoulder joint. In: *American Journal of Sports Medicine*. Vol 18. SAGE Publications; 1990:579-584. doi:10.1177/036354659001800604
11. Ward SR, Hentzen ER, Smallwood LH, et al. Rotator Cuff Muscle Architecture. *Clinical Orthopaedics and Related Research*. 2006;448(448):157-163. doi:10.1097/01.blo.0000194680.94882.d3
12. Halder AM, Zhao KD, O'Driscoll SW, Morrey BF, An KN. Dynamic contributions to superior shoulder stability. *Journal of*



*Orthopaedic Research*. 2001;19(2):206-212. doi:10.1016/S0736-0266(00)00028-0

13. Dugas JR, Campbell DA, Warren RF, Robie BH, Millett PJ. Anatomy and dimensions of rotator cuff insertions. *Journal of Shoulder and Elbow Surgery*. 2002;11(5):498-503. doi:10.1067/mse.2002.126208

14. Clark JM, Harryman DT. Tendons, ligaments, and capsule of the rotator cuff. Gross and microscopic anatomy. *Journal of Bone and Joint Surgery - Series A*. 1992;74(5):713-725. doi:10.2106/00004623-199274050-00010

15. Meister K. Injuries to the shoulder in the throwing athlete. Part one: Biomechanics/pathophysiology/classification of injury. *American Journal of Sports Medicine*. 2000;28(2):265-275. doi:10.1177/03635465000280022301

16. Lugo R, Kung P, Ma CB. Shoulder biomechanics. *European Journal of Radiology*. 2008;68(1):16-24. doi:10.1016/j.ejrad.2008.02.051

17. Dillman CJ, Fleisig GS, Andrews JR. Biomechanics of pitching with emphasis upon shoulder kinematics. *Journal of Orthopaedic and Sports Physical Therapy*. 1993;18(2):402-408. doi:10.2519/jospt.1993.18.2.402

18. Fortenbaugh D, Fleisig GS, Andrews JR. Baseball pitching biomechanics in relation to injury risk and performance. *Sports health*. 2009;1(4):314-320. doi:10.1177/1941738109338546

19. Serrano B, Serrano J. *Shoulder Injuries In Olympic Weightlifting: A Systematic Review*. Vol 2.; 2020. Accessed August 23, 2020. www.jmhsci.org

20. Garhammer J. Biomechanical Profiles of Olympic Weightlifters. *International Journal of Sport Biomechanics*. 2016;1(2):122-130. doi:10.1123/ijsb.1.2.122

21. Marius Viorel U, Vladimir P, Carmen O, Cosmina C. Biomechanical characteristics of movement phases of clean & jerk style in weightlifting performance. *Procedia - Social and Behavioral Sciences*. 2014;137:64-69. doi:10.1016/j.sbspro.2014.05.253

22. Grabe SA, Widule CJ. Comparative biomechanics of the jerk in olympic weightlifting. *Research Quarterly for Exercise and Sport*. 1988;59(1):1-8. doi:10.1080/02701367.1988.10605466

23. Blevins FT. Rotator cuff pathology in athletes. *Sports Medicine*. 1997;24(3):205-220. doi:10.2165/00007256-199724030-00009

24. Payne LZ, Altchek DW, Craig E v., Warren RF. Arthroscopic treatment of partial rotator cuff tears in young athletes. A preliminary report. *American Journal of Sports Medicine*. 1997;25(3):299-305. doi:10.1177/036354659702500305

25. Matava MJ, Purcell DB, Rudzki JR. Partial-thickness rotator cuff tears. *American Journal of Sports Medicine*. 2005;33(9):1405-1417. doi:10.1177/0363546505280213

26. Snyder SJ, Pachelli AF, del Pizzo W, Friedman MJ, Ferkel RD, Pattee G. Partial thickness rotator cuff tears: Results of arthroscopic treatment. *Arthroscopy: The Journal of Arthroscopic and Related Surgery*. 1991;7(1):1-7. doi:10.1016/0749-8063(91)90070-E

27. Romeo AA, Hang DW, Bach BR, Shott S. SECTION 11 ORIGINAL ARTICLES Repair of Full Thickness Rotator Cuff Tears Gender, Age, and Other Factors Affecting Outcome. Vol 367.

28. Hawkins RJ, Misamore GW, Hobeika PE. Surgery for full-thickness rotator-cuff tears. *Journal of Bone and Joint Surgery - Series A*. 1985;67(9):1349-1355. doi:10.2106/00004623-198567090-00007

29. Economopoulos KJ, Brockmeier SF. Rotator Cuff Tears in Overhead Athletes. *Clinics in Sports Medicine*. 2012;31(4):675-692. doi:10.1016/j.csm.2012.07.005

30. Costouros JG, Porratikul M, Lie DT, Warner JJP. Reversal of Suprascapular Neuropathy Following Arthroscopic Repair of Massive Supraspinatus and Infraspinatus Rotator Cuff Tears. *Arthroscopy - Journal of Arthroscopic and Related Surgery*. 2007;23(11):1152-1161. doi:10.1016/j.arthro.2007.06.014

31. Miller RM, Thunes J, Musahl V, Maiti S, Debski RE. Effects of tear size and location on predictions of supraspinatus tear propagation. *Journal of Biomechanics*. 2018;68:51-57. doi:10.1016/j.jbiomech.2017.12.017

32. Lyons AR, Tomlinson JE. Clinical diagnosis of tears of the rotator cuff. *Journal of Bone and Joint Surgery - Series B*. 1992;74(3):414-415. doi:10.1302/0301-620x.74b3.1587891

33. Jain NB, Wilcox RB, Katz JN, Higgins LD. Clinical Examination of the Rotator Cuff. *PM and R*. 2013;5(1):45-56. doi:10.1016/j.pmrj.2012.08.019

34. Hermans J, Luime JJ, Meuffels DE, Reijman M, Simel DL, Bierma-Zeinstra SMA. Does this patient with shoulder pain have rotator cuff disease? The rational clinical

examination systematic review. *JAMA - Journal of the American Medical Association*. 2013;310(8):837-847.

doi:10.1001/jama.2013.276187

35. Ouellette H, Labis J, Bredella M, Palmer WE, Sheah K, Torriani M. Spectrum of shoulder injuries in the baseball pitcher. *Skeletal Radiology*. 2008;37(6):491-498. doi:10.1007/s00256-007-0389-0

36. Fedoriw WW, Ramkumar P, McCulloch PC, Lintner DM. Return to play after treatment of superior labral tears in professional baseball players. In: *American Journal of Sports Medicine*. Vol 42. SAGE Publications Inc.; 2014:1155-1160.

doi:10.1177/0363546514528096

37. Fujii Y, Yoneda M, Wakitani S, Hayashida K. Histologic analysis of bony Bankart lesions in recurrent anterior instability of the shoulder. *Journal of Shoulder and Elbow Surgery*. 2006;15(2):218-223. doi:10.1016/j.jse.2005.06.006

38. Kim S-H, Ha K-I, Yoo J-C, Noh K-C. Kim's lesion: An incomplete and concealed avulsion of the posteroinferior labrum in posterior or multidirectional posteroinferior instability of the shoulder. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2004;20(7):712-720.

doi:10.1016/j.arthro.2004.06.012

39. Evancho AM, Stiles RG, Fajman WA, et al. MR imaging diagnosis of rotator cuff tears. *American Journal of Roentgenology*. 1988;151(4):751-754.

doi:10.2214/ajr.151.4.751

40. Zlatkin MB, Iannotti JP, Roberts MC, et al. Rotator cuff tears: Diagnostic performance of MR imaging. *Radiology*. 1989;172(1):223-229. doi:10.1148/radiology.172.1.2740508

41. Spielmann AL, Forster BB, Kokan P, Hawkins RH, Janzen DL. Shoulder after rotator cuff repair: MR imaging findings in asymptomatic individuals - Initial experience. *Radiology*. 1999;213(3):705-708. doi:10.1148/radiology.213.3.r99dc09705

42. Read JW, Perko M. Shoulder ultrasound: Diagnostic accuracy for impingement syndrome, rotator cuff tear, and biceps tendon pathology. *Journal of Shoulder and Elbow Surgery*. 1998;7(3):264-271. doi:10.1016/S1058-2746(98)90055-6

43. Charousset C, Bellaïche L, Duranthon LD, Grimberg J. Accuracy of CT arthrography in the assessment of tears of the rotator cuff. *The Journal of Bone and Joint Surgery British volume*. 2005;87-B(6):824-828. doi:10.1302/0301-620X.87B6.15836

44. Longo UG, Franceschi F, Berton A, Maffulli N, Droena V. Conservative Treatment and Rotator Cuff Tear Progression. In: *Medicine and Sport Science*. Vol 57. Karger Publishers; 2011:90-99.

doi:10.1159/000328910

45. Itoi E. Rotator cuff tear: Physical examination and conservative treatment. *Journal of Orthopaedic Science*. 2013;18(2):197-204. doi:10.1007/s00776-012-0345-2

46. Provencher CMT, Makani A, McNeil JW, Pomerantz ML, Golijanin P, Gross D. The Role of the Scapula in Throwing Disorders. *Sports Medicine and Arthroscopy Review*. 2014;22(2):80-87.

doi:10.1097/JSA.0000000000000023

47. Miyashita K, Kobayashi H, Koshida S, Urabe Y. Glenohumeral, scapular, and thoracic angles at maximum shoulder external rotation in throwing. *American Journal of Sports Medicine*. 2010;38(2):363-368.

doi:10.1177/0363546509347542

48. Vad VB, Gebeh A, Dines D, Altchek D, Norris B. Hip and shoulder internal rotation range of motion deficits in professional tennis players. *Journal of Science and Medicine in Sport*. 2003;6(1):71-75. doi:10.1016/S1440-2440(03)80010-5

49. Makaram NS, Safran MR, Abrams GD, Sherman SL, Murray IR. Rationale for the Use of Orthobiologics in Sports Medicine. *Operative Techniques in Sports Medicine*. Published online July 10, 2020:150753. doi:10.1016/j.otsm.2020.150753

50. Jang SJ, Kim J do, Cha SS. Platelet-rich plasma (PRP) injections as an effective treatment for early osteoarthritis. *European Journal of Orthopaedic Surgery and Traumatology*. 2013;23(5):573-580. doi:10.1007/s00590-012-1037-5

51. Kon E, Filardo G, di Martino A, Marcacci M. Platelet-rich plasma (PRP) to treat sports injuries: Evidence to support its use. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2011;19(4):516-527. doi:10.1007/s00167-010-1306-y

52. Mlynarek RA, Kuhn AW, Bedi A. *Platelet-Rich Plasma (PRP) in Orthopedic Sports Medicine*. Accessed August 23, 2020. www.amjorthopedics.com

53. Beck JJ, Murray MM, Christino MA. Clinical approach in youth sports medicine: Patients' and guardians' desired characteristics in sports medicine surgeons. *Journal of the American Academy of Orthopaedic Surgeons*.

2019;27(13):479-485. doi:10.5435/JAAOS-D-18-00263

54. Costa-Almeida R, Babo PS, Reis RL, Gomes ME. Platelet-rich Blood Derivatives for Tendon Regeneration. *The Journal of the American Academy of Orthopaedic Surgeons*. 2020;28(5):e202-e205. doi:10.5435/JAAOS-D-18-00638

55. Shams A, El-Sayed M, Gamal O, Ewes W. Subacromial injection of autologous platelet-rich plasma versus corticosteroid for the treatment of symptomatic partial rotator cuff tears. *European Journal of Orthopaedic Surgery and Traumatology*. 2016;26(8):837-842. doi:10.1007/s00590-016-1826-3

56. Hogan M v., Walker GN, Cui LR, Fu FH, Huard J. The role of stem cells and tissue engineering in orthopaedic sports medicine: Current evidence and future directions. *Arthroscopy - Journal of Arthroscopic and Related Surgery*. 2015;31(5):1017-1021. doi:10.1016/j.arthro.2014.11.033

57. Awad HA, Butler DL, Boivin GP, et al. Autologous mesenchymal stem cell-mediated repair of tendon. *Tissue Engineering*. 1999;5(3):267-277. doi:10.1089/ten.1999.5.267

58. Pas HIMFL, Moen MH, Haisma HJ, Winters M. No evidence for the use of stem cell therapy for tendon disorders: A systematic review. *British Journal of Sports Medicine*. 2017;51(13):996-1004. doi:10.1136/bjsports-2016-096794

59. Nourissat G, Diop A, Maurel N, et al. Mesenchymal Stem Cell Therapy Regenerates the Native Bone-Tendon Junction after Surgical Repair in a Degenerative Rat Model. Rannou FP, ed. *PLoS ONE*. 2010;5(8):e12248. doi:10.1371/journal.pone.0012248

60. Mora MV. Stem cell therapy in the management of shoulder rotator cuff disorders. *World Journal of Stem Cells*. 2015;7(4):691. doi:10.4252/wjsc.v7.i4.691

61. Jin H, Bae Y, Kim M, et al. Comparative Analysis of Human Mesenchymal Stem Cells from Bone Marrow, Adipose Tissue, and Umbilical Cord Blood as Sources of Cell Therapy. *International Journal of Molecular Sciences*. 2013;14(9):17986-18001. doi:10.3390/ijms140917986

62. Severud EL, Ruotolo C, Abbott DD, Nottage WM. All-arthroscopic versus mini-open rotator cuff repair: A long-term retrospective outcome comparison. *Arthroscopy - Journal of Arthroscopic and Related Surgery*.

2003;19(3):234-238.

doi:10.1053/jars.2003.50036

63. Burkhart SS, Danaceau SM, Pearce CE. Arthroscopic rotator cuff repair: Analysis of results by tear size and by repair technique-margin convergence versus direct tendon-to-bone repair. *Arthroscopy*. 2001;17(9):905-912. doi:10.1053/jars.2001.26821

64. van der Meijden OA, Westgard P, Chandler Z, Gaskill TR, Kokmeyer D, Millett PJ. Rehabilitation after arthroscopic rotator cuff repair: current concepts review and evidence-based guidelines. *International journal of sports physical therapy*. 2012;7(2):197-218. Accessed August 23, 2020. <http://www.ncbi.nlm.nih.gov/pubmed/22530194>

65. Düzgün I, Baltacı G, Ahmet Atay Ö. Comparison of slow and accelerated rehabilitation protocol after arthroscopic rotator cuff repair: Pain and functional activity. *Acta Orthopaedica et Traumatologica Turcica*. 2011;45(1):23-33. doi:10.3944/AOTT.2011.2386

66. Lee BG, Cho NS, Rhee YG. Effect of two rehabilitation protocols on range of motion and healing rates after arthroscopic rotator cuff repair: Aggressive versus limited early passive exercises. *Arthroscopy - Journal of Arthroscopic and Related Surgery*. 2012;28(1):34-42. doi:10.1016/j.arthro.2011.07.012